

Closed time like curves and the energy condition in 2+1 dimensional gravity

P. Menotti

Dipartimento di Fisica della Università, Pisa 56100, Italy and

INFN, Sezione di Pisa

D. Seminara

Scuola Normale Superiore, Pisa 56100, Italy and

INFN, Sezione di Pisa

Abstract

We consider gravity in 2+1 dimensions in presence of extended stationary sources with rotational symmetry. We prove by direct use of Einstein's equations that if i) the energy momentum tensor satisfies the weak energy condition, ii) the universe is open (conical at space infinity), iii) there are no CTC at space infinity, then there are no CTC at all.

The problem of closed time like curves (CTC) in 2+1 dimensional gravity has recently attracted considerable interest [1–11]. Gravity in 2+1 dimensions in addition to describing solutions of 3+1 dimensional gravity in presence of cosmic strings, is of interest in itself because due to its simplicity can be analyzed in great detail. Gott [1] was able to produce examples of kinematical configurations of point particles without spin which produce CTC in 2+1 dimensions. This system came under close scrutiny

[2–8]. Carrol, Fahri and Guth [4] prove that if the universe containing spinless point particles is open and has total time like momentum, no Gott pair can be created during its evolution thus supporting the idea that in an open timelike universe no CTC can form. 't Hooft [8] for a system of spinless point particles, using a general construction of a complete series of time ordered Cauchy surfaces, concludes that, also in the case of a closed universe, if Gott pairs are produced as envisaged in [4], the universe collapses before a CTC can form. Tipler [9] and Hawking [10] assuming the weak energy condition proved in 3+1 dimensions that if CTC are formed in a compact region of space-time, then one must necessarily have the creation of singularities. On the other hand it is very simple to recognize [2] that a point like source with angular momentum produce sufficiently near the source CTC. The existence of CTC appears to generate some irresolvable clash with causality [10]. In the case of the point like spinning source the appearance of CTC is usually ascribed to the unphysical nature of a point like spinning object [11]. A conjecture [2,10] states that if the metric is generated by a physical energy momentum tensor and proper boundary conditions are imposed, no CTC should develop. By physical sources one usually understands an energy momentum tensor which satisfies one or more among the weak , dominant or strong energy condition [12]. For this reason we investigate the problem in 2+1 dimensions in presence of extended sources in the simple instance of a stationary source with rotational symmetry. Our result is the following: if the energy momentum tensor satisfies the weak energy condition (WEC), the universe is open (conical at space infinity), and there are not CTC at space infinity, then there are no CTC at all. The proof follows from direct manipulation of Einstein' s equation combined with the WEC. The proof immediately extends to the case of cylindrical universes with angular momentum and to the case of “linear” universes.

In the treatment we shall extensively use the reduced radial gauge defined in [14] and

the method of solution developed in [15]. In stationary problems the reduced radial gauge [14] is defined by

$$\sum_i \xi^i \Gamma_{bi}^a = 0 \quad , \quad \sum_i \xi^i e_i^a = \sum_i \xi^i \delta_i^a \quad (1)$$

where the sums run over the space indices. (In the following the indices i, j, k, l, m run over space indices). In any space time dimensions there are resolvent formulae [14] which express the vierbeins and the connection in terms of the Riemann tensor in this gauge; they are analogous to those for the complete radial gauge [13]

$$\Gamma_{bi}^a(\boldsymbol{\xi}) = \xi^j \int_0^1 R_{bji}^a(\lambda \boldsymbol{\xi}) \lambda d\lambda \quad , \quad \Gamma_{b0}^a(\boldsymbol{\xi}) = \Gamma_{b0}^a(\mathbf{0}) + \xi^i \int_0^1 R_{bi0}^a(\lambda \boldsymbol{\xi}) d\lambda \quad (2)$$

$$e_i^a = \delta_i^a + \xi^l \xi^j \int_0^1 R_{jli}^a(\lambda \boldsymbol{\xi}) \lambda (1 - \lambda) d\lambda \quad , \quad e_0^a = \delta_0^a + \xi^i \Gamma_{i0}^a(\mathbf{0}) + \xi^i \xi^j \int_0^1 R_{ij0}^a(\lambda \boldsymbol{\xi}) (1 - \lambda) d\lambda \quad (3)$$

The simplifying feature of 2+1 dimensions is the substantial identification of the Riemann with the Ricci tensor and through Einstein's equation, with the energy momentum tensor. Explicitly, denoting with R^{ab} the Riemann two-form we have

$$R^{ab} = -\frac{\kappa}{4} \varepsilon^{abc} T_c = -\frac{\kappa}{8} \varepsilon^{abc} \varepsilon_{\rho\mu\nu} \tau_c{}^\rho dx^\mu \wedge dx^\nu, \quad (4)$$

where T_c is the energy momentum two form. Using such a relation one can express through a simple quadrature, the connections and the dreibeins in terms of the energy momentum tensor which is the source of the gravitational field and thus one solves Einstein's equation. On the other hand the energy momentum tensor is subjected to the covariant conservation law and symmetry condition. Thus our problem is to construct the general form of conserved symmetric sources in the reduced radial gauge which in addition should satisfy other physical requirements given by the support of the sources and the restrictions given by the energy condition [12].

The conservation and symmetry equations are

$$\mathcal{D}T^a = 0 \quad , \quad \varepsilon_{abc}T^b \wedge e^c = 0. \quad (5)$$

To solve these equations we use the technique developed in [15]. We introduce the cylindrical coordinates

$$\xi^0 = t \quad , \quad \xi^1 = \rho \cos \theta \quad , \quad \xi^2 = \rho \sin \theta \quad (6)$$

and the cotangent vectors

$$T_\mu = \frac{\partial \xi^0}{\partial \xi^\mu} \quad , \quad \Theta_\mu = \rho \frac{\partial \theta}{\partial \xi^\mu} \quad \text{and} \quad P_\mu = \frac{\partial \rho}{\partial \xi^\mu}. \quad (7)$$

In 2+1 dimensions the most general form of a radial $\Gamma_\mu^{ab}(\boldsymbol{\xi})$ is obviously given by

$$\Gamma_\mu^{ab}(\boldsymbol{\xi}) = \varepsilon^{abc} \varepsilon_{\mu\rho\nu} P^\rho A_c^\nu(\boldsymbol{\xi}) \quad (8)$$

and we parameterize $A_c^\nu(\boldsymbol{\xi})$ as follows

$$A_c^\rho(\boldsymbol{\xi}) = T_c \left[\beta_1 T^\rho + \frac{(\beta_2 - 1)}{\rho} \Theta^\rho \right] + \Theta_c \left[\alpha_1 T^\rho + \frac{\alpha_2}{\rho} \Theta^\rho \right] + P_c \left[\gamma_1 T^\rho + \frac{\gamma_2}{\rho} \Theta^\rho \right]. \quad (9)$$

We use for the coefficient of $T_c \Theta^\rho$ the form $\beta_2 - 1$ as it simplifies the writing of subsequent formulae. Using (9) and (3) we obtain for the dreibeins

$$e_0^a(\boldsymbol{\xi}) = -T^a A_1 - \Theta^a B_1 \quad , \quad e_i^a(\boldsymbol{\xi}) = -\frac{1}{\rho} \Theta^a \Theta_i B_2 - \frac{1}{\rho} T^a \Theta_i A_2 - P^a P_i \quad (10)$$

where A_i and B_i are defined by

$$A_1(\boldsymbol{\xi}) = \rho \int_0^1 \alpha_1(\lambda \boldsymbol{\xi}) d\lambda - 1 \quad , \quad A_2(\boldsymbol{\xi}) = \rho \int_0^1 \alpha_2(\lambda \boldsymbol{\xi}) d\lambda \quad , \quad B_i(\boldsymbol{\xi}) = \rho \int_0^1 \beta_i(\lambda \boldsymbol{\xi}) d\lambda \quad (11)$$

from which the metric is given by

$$ds^2 = (A_1^2 - B_1^2) dt^2 + 2(A_1 A_2 - B_1 B_2) dt d\theta + (A_2^2 - B_2^2) d\theta^2 - dr^2. \quad (12)$$

Substituting (9) in Einstein's equation (4) one obtains the most general energy momentum tensor in the radial gauge [14,15].

$$\begin{aligned} \tau^{a\rho} = & -\frac{4}{\kappa} \left\{ T^a \left(T^\rho \frac{\beta'_2}{\rho} + \Theta^\rho \beta'_1 \right) + \Theta^a \left(T^\rho \frac{\alpha'_2}{\rho} + \Theta^\rho \alpha'_1 \right) + P^a \left(T^\rho \frac{\gamma'_2}{\rho} + \Theta^\rho \gamma'_1 \right) + \right. \\ & \left. \frac{1}{\rho} P^\rho \left[T^a \left(\alpha_1 \gamma_2 - \alpha_2 \gamma_1 - \frac{\partial \beta_1}{\partial \theta} \right) + \Theta^a \left(\beta_1 \gamma_2 - \beta_2 \gamma_1 - \frac{\partial \alpha_1}{\partial \theta} \right) + P^a \left(\alpha_1 \beta_2 - \alpha_2 \beta_1 - \frac{\partial \gamma_1}{\partial \theta} \right) \right] \right\}. \end{aligned} \quad (13)$$

where the primes denote derivatives with respect to ρ . The vanishing of τ_c^ρ outside the source i.e. for $\rho \geq \rho_0$ implies that in such a region α_i , β_i , γ_i are constant in ρ and the coefficients of $P^\rho T_c$, $P^\rho \Theta_c$ and $P^\rho P_c$ vanish. In the case of rotational symmetry i.e. when α_i , β_i , γ_i do not depend on θ [15], it is easily shown that the symmetry condition on τ imposes $\gamma_i = 0$, and the contraction of $\tau^{a\mu}$ on the dreibein gives

$$\begin{aligned} \tau^{a\mu}(\xi) e_\mu^c(\xi) = \tau^{ac} = & -\frac{4}{\kappa \rho} \left\{ T^a \left[T^c \left(A_2 \beta'_1 - A_1 \beta'_2 \right) + \Theta^c \left(A_2 \alpha'_1 - A_1 \alpha'_2 \right) \right] + \right. \\ & \left. \Theta^a \left[T^c \left(B_2 \beta'_1 - B_1 \beta'_2 \right) + \Theta^c \left(B_2 \alpha'_1 - B_1 \alpha'_2 \right) \right] + P^a P^c \left(\alpha_1 \beta_2 - \alpha_2 \beta_1 \right) \right\}. \end{aligned} \quad (14)$$

The only surviving symmetry condition becomes

$$A_1 \alpha'_2 - A_2 \alpha'_1 + B_2 \beta'_1 - B_1 \beta'_2 = 0. \quad (15)$$

Eq. (15) is easily integrated; the regularity condition on $\Gamma_\mu^{ab}(\xi)$ at the origin imposes $\alpha_1 = o(\rho)$, $\alpha_2 = o(\rho)$, $\beta_1 = O(1)$ and $\beta_2 = 1 + o(\rho)$, which implies that the integration constant is 0 to get

$$A_1 \alpha_2 - A_2 \alpha_1 + B_2 \beta_1 - B_1 \beta_2 = 0. \quad (16)$$

It is immediately seen that a CTC implies the existence of a CTC with constant ρ and t . In fact if we call λ the parameter of the curve ($0 \leq \lambda < 1$), $t(\lambda)$ must satisfy $t(0) = t(1)$ and at the point λ_0 where $\frac{dt}{d\lambda} = 0$ we have that the tangent vector $(0, d\theta, d\rho)$ is time like i.e. also $(0, d\theta, 0)$ is time like, and thus the circle $t = 0$, $\rho = \rho(\lambda_0)$ is timelike. Thus to prove that CTC cannot exist is sufficient to prove that $g_{\theta\theta}$ which by assumption is negative at space infinity, cannot change sign. If the determinant of the dreibein in the radial gauge, where $g_{\rho\rho} \equiv -1$, vanishes for a

certain $\bar{\rho}$, it means that the manifold defined by $\rho = \text{const.}$ becomes one dimensional at $\rho = \bar{\rho}$, implying that at $\bar{\rho}$ the universe closes. Thus for an open universe we have that $\det(e) > 0$ for $\rho > 0$; in particular if we denote with ρ_0 a point outside the source we must have $\frac{d\det(e)}{d\rho} \big|_{\rho=\rho_0} \geq 0$ because for $\rho \geq \rho_0$, due to the support conditions, the determinant is a linear function of ρ for $\rho \geq \rho_0$. We show now that the WEC combined with $\det(e) > 0$ and the absence of CTC at infinity imply the absence of CTC for any ρ . In fact the WEC $v_a \tau^{ab} v_b \geq 0$ applied to the vectors $(1, 1, 0)$ and $(1, -1, 0)$ gives

$$\frac{d}{d\rho} [(\alpha_2 \pm \beta_2)(B_1 \pm A_1) - (B_2 \pm A_2)(\alpha_1 \pm \beta_1)] \geq 0 \quad (17)$$

which can be integrated between ρ and any point ρ_0 outside the source to give

$$E^{(\pm)}(\rho) \equiv (B_2 \pm A_2)(\alpha_1 \pm \beta_1) - (\alpha_2 \pm \beta_2)(B_1 \pm A_1) \geq (B_2^0 \pm A_2^0)(\alpha_1^0 \pm \beta_1^0) - (\alpha_2^0 \pm \beta_2^0)(B_1^0 \pm A_1^0) \quad (18)$$

We study now the sign of the r.h.s.. We have the following cases:

i) $(\alpha_2^0)^2 - (\beta_2^0)^2 \neq 0$ (conical universe).

Under the hypothesis that there are no CTC at infinity, as outside the source $g_{\theta\theta}$ becomes a quadratic polynomial in ρ , we have

$$(\alpha_2^0)^2 - (\beta_2^0)^2 < 0. \quad (19)$$

The support equation for $\tau^{\rho\rho}$ is

$$(\alpha_2^0 - \beta_2^0)(\alpha_1^0 + \beta_1^0) - (\alpha_2^0 + \beta_2^0)(\alpha_1^0 - \beta_1^0) = 0 \quad (20)$$

and using the equation of motion written in the form

$$\begin{aligned} (A_2 + B_2)(\alpha_1 - \beta_1) + (A_2 - B_2)(\alpha_1 + \beta_1) - \\ (\alpha_2 - \beta_2)(A_1 + B_1) - (\alpha_2 + \beta_2)(A_1 - B_1) = 0 \end{aligned} \quad (21)$$

and the determinant written in the form

$$\det(e) = \frac{1}{2}[(A_2 - B_2)(A_1 + B_1) - (A_2 + B_2)(A_1 - B_1)], \quad (22)$$

the r.h.s. of (18) becomes

$$E^{(\pm)}(\rho_0) = -\frac{\alpha_2^0 \pm \beta_2^0}{\alpha_2^0 \mp \beta_2^0} \frac{d}{d\rho} \det(e) \big|_{\rho=\rho_0} \quad (23)$$

which due to (19) and $\frac{d\det(e)}{d\rho} \big|_{\rho=\rho_0} \geq 0$ is non negative i.e. $E^{(\pm)}(\rho_0) \geq 0$.

ii) $\alpha_2 = \beta_2 \neq 0$ (linear universe).

Using the support equation we have $\alpha_1^0 = \beta_1^0$ which implies $E^{(-)}(\rho_0) = 0$. Using the equation of motion (21) and $\det(e) \neq 0$ one has $A_2^0 - B_2^0 \neq 0$ and thus we obtain

$$E^{(+)}(\rho_0) = -2 \frac{\alpha_2^0 + \beta_2^0}{A_2^0 - B_2^0} \det(e). \quad (24)$$

But $\det(e) > 0$ while the sign of $\frac{\alpha_2^0 + \beta_2^0}{A_2^0 - B_2^0}$ has to be negative if there are no CTC at infinity, because $A_2^2 - B_2^2$ in this case behaves linearly at infinity.

Similarly for $\alpha_2 = -\beta_2$ we obtain $E^{(+)}(\rho_0) = 0$ and $E^{(-)}(\rho_0) > 0$.

iii) If $\alpha_1^0 = \alpha_2^0 = \beta_1^0 = \beta_2^0 = 0$ (cylindrical universe with angular momentum) we have also $E^{(\pm)}(\rho_0) = 0$.

The only case which escapes our analysis is $\alpha_2^0 = \beta_2^0 = 0$ and α_1^0 and/or $\beta_1^0 \neq 0$; this situation corresponds to a cylindrical universe generated by a string with tension and total angular momentum 0. Thus except for this case we have that the r.h.s. of equation (18) is $E^{(\pm)}(\rho_0) \geq 0$. Let us now consider the following combination

$$(A_2 - B_2)^2 E^{(+)}(\rho) + (A_2 + B_2)^2 E^{(-)}(\rho) \geq 0. \quad (25)$$

A little algebra shows that the l.h.s. equals

$$-2\det^2(e) \frac{d}{d\rho} \left(\frac{A_2^2(\rho) - B_2^2(\rho)}{\det(e)} \right). \quad (26)$$

Thus we reached the conclusion that $\frac{d}{d\rho} \left(\frac{A_2^2(\rho) - B_2^2(\rho)}{\det(e)} \right) \leq 0$; which means that $\left(\frac{A_2^2(\rho) - B_2^2(\rho)}{\det(e)} \right)$ is a non increasing function of ρ . As $\left(\frac{A_2^2(\rho) - B_2^2(\rho)}{\det(e)} \right)$ at the origin is zero we obtain that $A_2^2(\rho) - B_2^2(\rho)$ is always negative and thus we cannot have CTC.

It is not difficult to produce examples in which all energy conditions are satisfied and CTC exist for any radius larger than a given one; but we obviously violate the condition that no CTC exists at infinity. Also renouncing to the energy conditions one can produce examples [14] in which non CTC exist at infinity, but they exist at finite radius. As mentioned above the only case of open universe with no CTC at infinity which escapes our analysis is the rather unphysical situation of a cylindrical universe with zero angular momentum of the type generated by a closed string with tension [16] for which on the other hand no CTC exists outside the source. We have no example of an extended source satisfying the energy condition and producing such a universe.

We mention that the approach with the radial gauge, for which the solution can be given in terms of quadratures [14,15], works also for the case of rotational symmetry with time dependent sources [15] and for stationary problems in absence of rotational symmetry [17]; one could apply them to the problem of CTC in 2+1 dimensions in absence of rotational symmetry and in presence of time dependence.

REFERENCES

- [1] J.R. Gott, *Phys. Rev. D* 46 (1991) 1126.
- [2] S. Deser, R. Jackiw and G. 't Hooft, *Phys. Rev. Lett.* 68 (1992) 2647.
- [3] S.M. Carrol, E. Farhi and A.H. Guth, *Phys. Rev. Lett.* 68 (1992) 263; Erratum: 68 (1992) 3368.
- [4] S.M. Carrol, E. Farhi and A.H. Guth, “ *Gott time machines cannot exist in an open (2+1) dimensional universe with timelike total momentum*” preprint MIT-CTP-2117 (1992)
- [5] C. Cutler, *Phys. Rev. D* 45 (1992) 487.
- [6] D. Kabat, *Phys. Rev. D* 46 (1992) 2720.
- [7] A. Ori, *Phys. Rev. D* 44 (1992) R2214.
- [8] G. 't Hooft, *Class. Quantum Grav.* 9 (1992) 1335.
- [9] F. J. Tipler, *Ann. Phys. (N.Y.)* 108 (1977) 1.
- [10] S. W. Hawking, *Phys. Rev. D* 46 (1992) 603.
- [11] S. Deser and R. Jackiw, *Time Travel* preprint BRX TH 334; CTP#2101 (1992).
- [12] S.W. Hawking and G.F.R. Ellis, *The large scale structure of space-time*, p. 33, Cambridge University Press, Cambridge, 1976.
- [13] G. Modanese and M. Toller, *J. Math. Phys.* 31 (1990) 452.
- [14] P. Menotti and D. Seminara, *Ann. Phys. (N.Y.)* 208 (1991) 449.
- [15] P. Menotti and D. Seminara, *Nucl. Phys.* B376 (1992) 411.

- [16] S. Deser and B. Laurent, *Gen. Relativ. Gravitation* 18 (1986) 617; G. Clement
Ann. Phys. (N.Y.) 201 (1990) 241.
- [17] P. Menotti and D. Seminara, to be published.